

R Textbook Companion for
Statistics - Concepts and Applications
by Nabendu Pal and Sahadeb Sarkar¹

Created by
M. Sai Anand
Bachelor of Technology
Computer Science and Engineering
Kalasalingam Academy of Research and Education
Cross-Checked by
R TBC Team

June 8, 2021

¹Funded by a grant from the National Mission on Education through ICT
- <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and R
codes written in it can be downloaded from the "Textbook Companion Project"
section at the website - <https://r.fossee.in>.

Book Description

Title: Statistics - Concepts and Applications

Author: Nabendu Pal and Sahadeb Sarkar

Publisher: Phi Learning Private Limited, Sonapat, Harayana.

Edition: 2

Year: 2013

ISBN: 9788120334458

R numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means an R code whose theory is explained in Section 2.3 of the book.

Contents

List of R Codes	4
2 Organizing a Raw Dataset	5
3 Pictorial Representation of a Dataset	10
4 Summarizing a Raw Dataset	12
5 Summarizing an Organized Dataset	20
6 Concepts of Probability	23
7 Random Variables	34
8 Binomial Experiments	40
9 Normal Curve and Normal Distribution	44
10 Applications of the Normal Distribution	47
11 Estimation of Population Parameters	52
12 Hypothesis Testing for a Single Population	55
13 Hypothesis Testing to Compare Two Populations	60
14 Bivariate Quantitative Data Correlation and Regression	65
15 Bivariate Catagorical Data Contingency Table	69

16	Multinomial Experiments Goodness of Fit Test	73
17	Hypothesis Testing to Compare Multiple Populations	78
18	Quality Management Using Statistics	82

List of R Codes

Exa 2.1	Relative frequency of dataset on voters preference . . .	5
Exa 2.2	Relative frequency of dataset on gender painkiller study	6
Exa 2.3	Relative frequency of dataset on number of individuals treated at ER per day	6
Exa 2.4	Relative frequency of dataset on for the oak tree circum- ference	8
Exa 3.1	Stem leaf display of commuting distance data	10
Exa 3.2	Time plot of burglaries between 1981 and 1990	10
Exa 4.1	Computation of mode for distinct entry	12
Exa 4.2	Computation of mode for dataset	12
Exa 4.3	Median of the given dataset	13
Exa 4.4	Median of the given dataset	13
Exa 4.5	Median of the 2 datasets	14
Exa 4.6	Median of the 2 datasets	14
Exa 4.7	Computation of mean for the dataset	15
Exa 4.8.i	Construction of datasets with same medians but differ- ent means	15
Exa 4.8.ii	Construction of datasets with same means but different medians	15
Exa 4.8.iii	Construction of datasets with same medians but differ- ent modes	16
Exa 4.8.iv	Construction of datasets with same means medians and modes	16
Exa 4.9	Range length of elementary school children dataset . . .	17
Exa 4.10	Range and IQR of given dataset	17
Exa 4.11	Range and IQR of given dataset	18
Exa 4.12	Variance of the given dataset	18

Exa 4.13	Box whisker plots for males and females working in tele- com sector	19
Exa 5.1	Computation of central tendency and variability for the grouped dataset on the number of patients treated per day at the emergency room	20
Exa 5.2	Computation of variability for the randomly generated grouped dataset with the given class and frequency on the population of all working mothers in a large city	21
Exa 6.1	Probability of getting head in tossing a fair coin	23
Exa 6.3	Probability of getting face value less than 3 in rolling a die	23
Exa 6.6	Probability of getting a queen card from a deck of 52 cards	24
Exa 6.7	Probability of getting 2 heads in tossing a fair coin twice in a row	24
Exa 6.8	Probability of getting 2 heads in tossing a fair coin thrice in a row	24
Exa 6.9.a	Probability of getting red marbles from a box of red and blue marbles when there are 30 red and 40 blue marbles	25
Exa 6.9.b	Probability of getting red marbles from a box of red and blue marbles when composition of box is unknown	25
Exa 6.10	Getting 4 or more face values from the unbalanced die with known sample points	26
Exa 6.11.a	Probability of students failed in maths test from a class of 120 students	26
Exa 6.11.b	Probability of students passed in both test from a class of 120 students	27
Exa 6.11.c	Probability of students failed in both test from a class of 120 students	27
Exa 6.12	Probability of getting a card that is either red or a king from a deck of 52 cards	28
Exa 6.13	Probability of getting atleast 1 head in tossing a fair coin four times in a row	29
Exa 6.14	Probability of drawing a card that is King and knowing that there is no Queen in deck of 52 cards	29
Exa 6.15	Probability of a student passed in maths test given that he or she is passed in atleast 1 test from a class of 120 students	30

Exa 6.16	Independent events of cards being drawn at random that are either red or a king from deck of 52 cards	30
Exa 6.17	Probability of a ship departing on time	31
Exa 6.18	Probability of getting a blue marble during first draw given that the second draw gave a blue marble from 2 boxes using bayes theorem	32
Exa 7.1	Random variable operations on a rolling die	34
Exa 7.2	Random variable operations on tossing a coin twice	35
Exa 7.3	Random variable operations in tossing a fair coin twice and then rolling a balanced die	36
Exa 7.4	Random variable operations on a sample of 3 voters	37
Exa 7.6	To find all possible grade pairs for a certain mark obtained	38
Exa 8.1	Probability of answering correctly in a pop quiz	40
Exa 8.2	Probability of finding mortar shell explosion on enemy target	41
Exa 8.3	Probability of at least 1 sensor detecting gas leak	41
Exa 8.4	Probability of fire accident due to arson	42
Exa 8.5	Probability of members who are seeking capital punishment	42
Exa 8.6	Tossing a fair coin 15 times	42
Exa 9.1	Percentage of children facing difficulties with new teaching method	44
Exa 9.2.a	Probability for a standard normal random variable	44
Exa 9.2.b	Probability for a standard normal random variable	45
Exa 9.2.c	Probability for a standard normal random variable	45
Exa 9.2.d	Probability for a standard normal random variable	45
Exa 9.3	Percentage of day time temperature between 80 to 100	46
Exa 10.1	Probability of 6 or more people suffering from hypothermia	47
Exa 10.2	Probability of 80 or less people suffering from hypothermia	47
Exa 10.3.a	Probability of 640 or more students passing the exam	48
Exa 10.3.b	Probability of 600 or fewer students passing the exam	48
Exa 10.3.c	Probability of students passing the exam between 585 and 625	49
Exa 10.4	Probability of sample batteries having average life of 52 or less	49

Exa 10.5.a	Distribution of sample average of bacteria count per ML of milk	50
Exa 10.5.b	Probability of sample having bacteria between 2400 to 2700	50
Exa 10.6	Probability of sample average of IQ scores not differing from population mean by more than 1 point	51
Exa 11.2	Computing a 90 percent confidence interval for mean PE ratio of all utility stocks in NYSE	52
Exa 11.3	Computing 95 percent confidence interval for the mean January average temperature in Phoenix	52
Exa 11.4	Computing 90 percent confidence interval for the percentage of US parents whose kids favourite food is pizza	53
Exa 12.2	Hypothesis testing on claim of mean household income in a business district	55
Exa 12.3	Hypothesis testing on claims of nutritionist on the rise of potato consumption	55
Exa 12.4	Hypothesis testing on claims based on students councils on time spent in library	56
Exa 12.5	Hypothesis testing on claims of management on share of fruits is 50 percent	57
Exa 12.6	Hypothesis testing on claims on weight of the newly minted coins	57
Exa 12.7	Verifying Normality for the given dataset	58
Exa 13.2	Testing the claims on painkiller which are identical in terms of mean time required for body absorption	60
Exa 13.3	Testing the difference between the mean stopping distances of two types of tyres	61
Exa 13.4	Testing on effects of a new hypertension drug which changes the body temperature	62
Exa 13.5	Testing the beliefs on the methods which is superior than other in treating patients with broken arm	62
Exa 13.6	Testing on data on ball bearing from 2 machines indicating that the machines are identical	63
Exa 14.1	Computing the operations of correlation and regression on population consisting of all married couples	65
Exa 14.2	Computing the operations of correlation and regression on population of sales of hot chocolate and temperature for a snack shop	67

Exa 15.1	Testing independence of gender wise usage of 2 types of painkillers	69
Exa 15.2	Testing independence on whether smoking causes hypertension on dataset of randomly selected set of patients	70
Exa 15.3	Testing independence of accidents involved in accordance with BAC levels and the region	71
Exa 16.1	Correlation coefficient describing the probability distribution of the outcome when 15 marbles are drawn at random from a box of 100 where 60 is red and 40 is blue	73
Exa 16.3	Finding the probability of rolling a die and getting multiple faces	73
Exa 16.4	Finding the probability of getting five 6s by rolling a die 15 times	74
Exa 16.5	Testing the hypothesis of having 60 percent red marbles from a box having unknown composition of red and blue marbles	74
Exa 16.6	Determining the uniformity in composition of types of marble in a box	75
Exa 16.7	Testing the hypothesis of ratio of the offsprings in a biology experiment	75
Exa 16.8	Goodness of fit test of survey data on a city daytime traffic	76
Exa 17.1	Computing the population mean variance and applying 1 way anova test on experimental data of number of seconds taken to help in presence of other persons . .	78
Exa 17.2	Classification of two way ANOVA and Barlett's test on data studying weight gains by rats influenced by the source of protein	79
Exa 18.1	To test whether the manufacturing process is in control by construction of x chart and s chart from the dataset of JetMate	82
Exa 18.2	To test whether the bottling process is in control by construction of x chart and r chart and s chart from the dataset of BottlePack company	83
Exa 18.3	To test whether the manufacturing process is in control by construction of s chart from the dataset of Blackstone company	84

Exa 18.4	To test whether the manufacturing process is in control by construction of p chart from the dataset of ChipIt- Good company	84
----------	---	----

Chapter 2

Organizing a Raw Dataset

R code Exa 2.1 Relative frequency of dataset on voters preference

```
1 #Page Number: 9
2
3 dataset<- read.csv("dataset.csv", header = T)
4 print(dataset)
5 A <- dataset[ dataset$Vote == 'A', ]
6 AF_A <-length(A$Population)
7 B <- dataset[ dataset$Vote == 'B', ]
8 AF_B <-length(B$Population)
9 C <- dataset[ dataset$Vote == 'C', ]
10 AF_C <-length(C$Population)
11 Total_Freq <- length(dataset$Population)
12 RF_A <- AF_A / Total_Freq
13 print(RF_A)
14 RF_B <- AF_B / Total_Freq
15 print(RF_B)
16 RF_C <- AF_C / Total_Freq
17 print(RF_C)
18 Total_RF = RF_A+RF_B+RF_C
19 print(Total_RF)
```

R code Exa 2.2 Relative frequency of dataset on gender painkiller study

```
1 #Page Number: 10
2
3 Tot_F = 0.60 * 500
4 Tot_M = 500 - Tot_F
5 F1 = 0.45 * 300
6 F2 = 300 - F1
7 MED_1 = 0.50 * 500
8 MED_2 = 500 - MED_1
9 M1 = MED_1 - F1
10 M2 = Tot_M - M1
11 req <- data.frame(GENDER = c("M", "F"), Medicine_1 =
      c(M1, F1), Medicine_2 = c(M2, F2))
12 RF_M1 = M1 / 500
13 RF_M2 = M2 / 500
14 RF_F1 = F1 / 500
15 RF_F2 = F2 / 500
16 genfreq <- data.frame( Catagory = c('M1', 'M2', 'F1', '
      F2' ), Frequency = c(M1,M2,F1,F2), Relative_
      Frequency = c(RF_M1,RF_M2,RF_F1,RF_F2) )
17 print(genfreq)
18 cat(paste(" Total Frequency:", sum(genfreq$Frequency),
      "\nTotal Relative Frequency:", sum(genfreq$
      Relative_Frequency)))
```

R code Exa 2.3 Relative frequency of dataset on number of individuals treated at ER per day

```
1 #Page Number: 12
2
3 dataset <- read.csv("data.csv", header = T)
```

```

4
5 minc1 <- min(dataset$C1)
6 minc2 <- min(dataset$C2)
7 minc3 <- min(dataset$C3)
8 minc4 <- min(dataset$C4)
9 minc5 <- min(dataset$C5)
10 minc6 <- min(dataset$C6)
11 minc7 <- min(dataset$C7)
12 minc8 <- min(dataset$C8)
13
14
15 maxc1 <- max(dataset$C1)
16 maxc2 <- max(dataset$C2)
17 maxc3 <- max(dataset$C3)
18 maxc4 <- max(dataset$C4)
19 maxc5 <- max(dataset$C5)
20 maxc6 <- max(dataset$C6)
21 maxc7 <- max(dataset$C7)
22 maxc8 <- max(dataset$C8)
23
24 Column_Min = c(minc1,minc2,minc3,minc4,minc5,minc6,
    minc7,minc8)
25 print(Column_Min)
26
27 Column_Max = c(maxc1,maxc2,maxc3,maxc4,maxc5,maxc6,
    maxc7,maxc8)
28 print(Column_Max)
29
30 Up_Lim <- max(dataset) + 0.5
31 Low_Lim <- min(dataset) - 0.5
32 L <- Up_Lim - Low_Lim
33
34
35 k <- 5
36 l <- L/k
37
38
39 breaks = seq( Low_Lim, Up_Lim, by= l )

```

```

40
41
42 frequency.cut<- cut(x<- c
      (40,8,26,21,36,35,42,32,46,30,42,53,28,22,40,6,17,28,19,38,13,25,
      ,breaks)
43
44 dataset.freq <- table(frequency.cut)
45
46 cbind(dataset.freq)
47
48
49 Relative_Freq <- dataset.freq/ 40
50 print(Relative_Freq)
51 cat(" Total Frequency:",sum(dataset.freq)," \nTotal
      Relative Frequency:",sum(Relative_Freq))

```

R code Exa 2.4 Relative frequency of dataset on for the oak tree circumference

```

1 #Page Number: 13
2
3 circum_table <- data.frame(Circumference = c('10-24'
      , '25-39', '40-54', '55-69' ), No._Of_Tree = c
      (8,22,53,17))
4 mod_class_table <- data.frame(Class_modified = c('
      9.5-24.5', '24.5-39.5', '39.5-54.5', '54.5-69.5')
      )
5 REL_Freq <- circum_table$No._Of_Tree / 100
6 Modfreq_Table <- data.frame(mod_class_table,
      Frequency = circum_table$No._Of_Tree, Relative_
      Frequency = REL_Freq)
7 final <- data.frame(Class = mod_class_table,
      Frequency = circum_table$No._Of_Tree, Rel.
      Frequency = REL_Freq)
8 print(final)

```

```
9 cat("Total Frequency:",sum(circum_table$No._Of_Tree)
    ,"\nTotal Relative Frequency:",sum(REL_Freq))
```

Chapter 3

Pictorial Representation of a Dataset

R code Exa 3.1 Stem leaf display of commuting distance data

```
1 #Page Number: 33
2
3 data <- data.frame(dist = c
  (13,7,12,6,34,14,47,25,45,2,13,26,10,8,1,14,41,10,3,21,8,13,28,24
  )
4 stem(data$dist)
```

R code Exa 3.2 Time plot of burglaries between 1981 and 1990

```
1 #Page Number: 34
2
3 library(ggplot2)
4 data <- data.frame(YEAR = c('1981', '1982', '1983', '
  1984', '1985', '1986', '1987', '1988', '1989', '1990'),
  Theft = c(24,18,16,27,30,34,37,38,40,39) )
5 ggplot(data, aes(YEAR, Theft, group = 1)) +
```

```
6 geom_point(shape = 0) +  
7 geom_line()+  
8 xlab("Year")+ ylab("Theft")
```

Chapter 4

Summarizing a Raw Dataset

R code Exa 4.1 Computation of mode for distinct entry

```
1 #Page Number: 55
2
3 library(plyr)
4
5 dataset <- data.frame(Distinct_entry = c
6   (8,9,0,9,1,4,3,7))
7
8
9 y <- count(dataset, 'Distinct_entry')
10 print(y)
11
12
13 z <- table(dataset)
14 cat("Mode is", names(z)[which(z==max(z))] )
```

R code Exa 4.2 Computation of mode for dataset

```

1 #Page Number: 56
2
3 library(plyr)
4
5 dataset <- data.frame(Distinct_entry = c
      (-1,-2,3,0,3,-1,4,2,7,5))
6 n <- length(dataset$Distinct_entry)
7
8
9 y <- count(dataset, 'Distinct_entry')
10 print(y)
11
12
13 z <- table(dataset)
14 cat("Mode is", names(z)[which(z==max(z))] )

```

R code Exa 4.3 Median of the given dataset

```

1 #Page Number:56
2
3 data <- c(2,-1,0,4,7,7,8,0,11)
4
5
6 ascend_order <- sort(data)
7
8
9 M <- median(ascend_order)
10 print(M)

```

R code Exa 4.4 Median of the given dataset

```

1 #Page Number: 56
2

```

```
3 data <- c(4,7,9,1,3,2,1,1,7,0)
4
5
6 ascend_order <- sort(data)
7
8 M <- median(ascend_order)
9 print(M)
```

R code Exa 4.5 Median of the 2 datasets

```
1 #Page Number: 56
2
3 data1 <- c(0,1,1,1,10)
4 data2 <- c(-27,1,1,2,1000)
5
6
7 ascend_order1 <- sort(data1)
8 ascend_order2 <- sort(data2)
9
10
11 M1 <- median(ascend_order1)
12 M2 <- median(ascend_order2)
13 print(M1)
14 print(M2)
```

R code Exa 4.6 Median of the 2 datasets

```
1 #Page Number: 56
2
3 data1 <- c(0,1,1,1,1,1,2)
4 data2 <- c(-100,-100,-100,1,100,100,200)
5
6
```

```
7 ascend_order1 <- sort(data1)
8 ascend_order2 <- sort(data2)
9
10
11 M1 <- median(ascend_order1)
12 M2 <- median(ascend_order2)
13 print(M1)
14 print(M2)
```

R code Exa 4.7 Computation of mean for the dataset

```
1 #Page Number : 57
2
3 data <- c(5,-1,0,2,4)
4 mean(data)
```

R code Exa 4.8.i Construction of datasets with same medians but different means

```
1 #Page Number: 57
2
3 dataset1 <- c(5,6,6,9,9)
4 dataset2 <- c(3,6,6,6,9)
5
6 cat("Mean of dataset 1:",mean(dataset1)," \nMean of
  dataset 2:",mean(dataset2))
7 cat("Median of dataset 1:",median(dataset1)," \nMedian of dataset 2:",median(dataset2))
```

R code Exa 4.8.ii Construction of datasets with same means but different medians

```

1 #Page Number: 57
2
3 dataset1 <- c(1,1,3,9,11)
4 dataset2 <- c(1,1,1,11,11)
5
6
7 cat("Mean of dataset 1:",mean(dataset1),"\\nMean of
  dataset 2:",mean(dataset2))
8 cat("Median of dataset 1:",median(dataset1),"\\
  nMedian of dataset 2:",median(dataset2))

```

R code Exa 4.8.iii Construction of datasets with same medians but different modes

```

1 #Page Number: 57
2
3 dataset1 <- c(1,1,1,2,2)
4 dataset2 <- c(1,1,3,4,5)
5
6
7 cat("Median of dataset 1:",median(dataset1),"\\
  nMedian of dataset 2:",median(dataset2))
8
9
10 x <- table(dataset1)
11 y <- table(dataset2)
12
13 cat("Mode of dataset 1:",names(x)[which(x==max(x))],
  "\\nMode of dataset 2:",names(y)[which(y==max(y))
  ])

```

R code Exa 4.8.iv Construction of datasets with same means medians and modes

```

1 #Page Number: 57
2
3 dataset1 <- c(1,2,2,2,3)
4 dataset2 <- c(0,2,2,2,4)
5
6
7 x <- table(dataset1)
8 cat("Mean:",mean(dataset1),"Median:",median(dataset1
9     ),"Mode:",as.numeric(names(x)[which(x==max(x))]))
10
11 y <- table(dataset2)
12 cat("Mean:",mean(dataset2),"Median:",median(dataset2
13     ),"Mode:",as.numeric(names(y)[which(y==max(y))]))

```

R code Exa 4.9 Range length of elementary school children dataset

```

1 #Page Number: 57
2
3 data <- c(72,113,97,91,105,88,119)
4
5 x <- min(data)
6 y <- max(data)
7
8 RL <- y-x
9 print(paste('The Range Length is:', RL))

```

R code Exa 4.10 Range and IQR of given dataset

```

1 #Page Number: 58
2
3 data <- c(5,7,-1,0,2,9,9,4,3,11)
4
5 x <- min(data)

```



```
6 y <- max(data)
7 cat('The range is:',x,"",y)
8
9 IQR_data <- quantile(data,type = 2)
10 cat('The IQR is:',IQR_data[2],"",IQR_data[4] )
```

R code Exa 4.11 Range and IQR of given dataset

```
1 #Page Number: 58
2
3 data <- c(5,7,-1,0,2,9,9,4,3,11,6)
4
5 x <- min(data)
6 y <- max(data)
7 RL <- y - x
8 print(paste('The range is:', RL))
9
10
11 IQR_data <- IQR(sort(data), type = 1)
12 print(paste('The IQRL is:',IQR_data ))
```

R code Exa 4.12 Variance of the given dataset

```
1 #Page Number: 59
2
3 data <- c(-2,3,0,2,2)
4
5 y <- var(data)
6 print(paste('The Variance is :',y ))
```

R code Exa 4.13 Box whisker plots for males and females working in tele-
com sector

```
1 #Page Number: 60
2
3 data <- read.csv("data.csv", header = T)
4
5
6 boxplot(data$MALES,main = "Box-whisker plot for Male
      ", ylim = c(10,80) )
7 boxplot(data$FEMALES,main = "Box-whisker plot for
      Female", ylim = c(10,80) )
```

Chapter 5

Summarizing an Organized Dataset

R code Exa 5.1 Computation of central tendency and variability for the grouped dataset on the number of patients treated per day at the emergency room

```
1 #Page Number: 68
2
3 data <- c
      (40,8,26,21,36,35,42,32,46,30,42,53,28,22,40,6,17,28,19,38,13,25,
4 n <- length(data)
5 interval <- seq(5.5, 60.5 , by = 11)
6 class <- cut(data, interval)
7 frequency <- cbind(table(class))
8
9 RF <- frequency / n
10 CRF <- cumsum(frequency) / n
11
12
13 midpoints <- function(x, dp=2){
14   lower <- as.numeric(gsub(" ,.*", "", gsub("
      \\(\\|\\|\\|\\|\\|\\|", "", x)))
```

```

15   upper <- as.numeric(gsub(".", "", gsub(
      "\\(|\\|\\|\\|\\|\\|)", "", x)))
16   return(round(lower+(upper-lower)/2, dp))
17 }
18
19 Midpoint <- sort(unique(midpoints(class)))
20 Midfreq <- Midpoint * RF
21 xbar <- sum(Midfreq)
22 k <- sum((Midpoint - xbar)^2 * frequency)
23 variance <- k / (n-1)
24 sd <- sqrt(variance)
25 cat(" Variance :", round(variance, 2), "& Standard
      Deviation:", round(sd, 2))
26
27 Q1 <- interval[2] + ((Midpoint[1] / RF[2]) * (0.25 -
      CRF[1] ))
28 Q2 <- interval[3] + ((Midpoint[1] / RF[3]) * (0.5 -
      CRF[2] ))
29 Q3 <- interval[3] + ((Midpoint[1] / RF[3]) * (0.75 -
      CRF[2] ))
30 cat(" Q1=", round(Q1, 1), " Q2=", round(Q2, 2), " Q3=", round(
      Q3, 2))

```

R code Exa 5.2 Computation of variability for the randomly generated grouped dataset with the given class and frequency on the population of all working mothers in a large city

```

1 #Page Number: 72
2
3 random_input_data <- c
      (6,11,21,39,45,9,18,29,36,41,2,18,27,7,19,21,1,13,29,4,20,28,7,1,
4 n <- length(random_input_data)
5
6 interval <- seq( from = 0.5, to = 50.5, by = 10)

```

```

7 class <- cut(random_input_data, interval)
8 frequency <- cbind(table(class))
9
10
11 RF <- frequency / n
12
13
14
15 midpoints <- function(x, dp=2){
16   lower <- as.numeric(gsub(",.*", "", gsub("
17     \\(\\|\\|\\|\\|) \\|\\|", "", x)))
18   upper <- as.numeric(gsub(".*", "", gsub("
19     \\(\\|\\|\\|\\|) \\|\\|", "", x)))
20   return(round(lower+(upper-lower)/2, dp))
21 }
22
23 a <-midpoints(class)
24
25 Midpoint <- unique(a)
26
27 Mid_Freq <- Midpoint * frequency
28
29 b <- sum(Mid_Freq)
30
31 m <- b/n
32
33 S <- (Midpoint - m)^2 * frequency
34 Var <- (sum(S)) / (n-1)
35 print(paste('Variance is:', round(Var,2)))
36 SD <- sqrt(Var)
37 print(paste('Standard Deviation is:', round(SD,2)))

```

Chapter 6

Concepts of Probability

R code Exa 6.1 Probability of getting head in tossing a fair coin

```
1 #Page Number: 81 & 83(cont.)
2
3 library(prob)
4 a <- tosscoin(1)
5 c <- length(a[a$toss1 == 'H']) / length(a$toss1)
6 print(paste('The probability is :',c))
```

R code Exa 6.3 Probability of getting face value less than 3 in rolling a die

```
1 #Page Number: 83
2
3 library(prob)
4 a <- rolldie(1)
5 c <- length(a[a$X1 < 3,]) / length(a$X1)
6 print(round(c,4))
```

R code Exa 6.6 Probability of getting a queen card from a deck of 52 cards

```
1 #Page Number: 83
2
3 library(prob)
4 a <- cards()
5 b <- subset(a, rank == 'Q')
6 c <- length(b$rank) /length(a$rank)
7 print(paste(' The probability is :',round(c,3)))
```

R code Exa 6.7 Probability of getting 2 heads in tossing a fair coin twice in a row

```
1 #Page Number: 83
2
3 library(prob)
4 S <- tosscoin(2,makespace = T)
5 P <- S[1,3]
6 print(P)
```

R code Exa 6.8 Probability of getting 2 heads in tossing a fair coin thrice in a row

```
1 #Page Number: 83
2
3 library(prob)
4
5 S <- tosscoin(3, makespace = T)
6 H1 <- S[2, 'probs']
7 H2 <- S[3, 'probs']
8 H3 <- S[5, 'probs']
9 Pro <- H1+H2+H3
10 print(paste('The probability is:',Pro))
```

R code Exa 6.9.a Probability of getting red marbles from a box of red and blue marbles when there are 30 red and 40 blue marbles

```
1 #Page Number: 84
2
3 library(prob)
4 S <- rep(c("Red", "Blue"), times = c(30,40))
5 P <- urnsamples(S, size = 1, replace = F, ordered =
  F)
6
7
8 R <- P[P$out == 'Red',]
9
10 P_R = length(R) / length(P$out)
11 print(round(P_R,4))
12
13 #The answer provided in the textbook is wrong.
```

R code Exa 6.9.b Probability of getting red marbles from a box of red and blue marbles when composition of box is unknown

```
1 #Page Number: 84
2
3 unknown_comp <- c(3,3,8,24.5,53,112,165,226)
4 n <- c(5,10,20,50,100,200,300,400)
5
6 r_n <- unknown_comp / n
7 print(r_n)
```

R code Exa 6.10 Getting 4 or more face values from the unbalanced die with known sample points

```
1 #Page Number: 85
2
3 library(prob)
4 library(MASS)
5 S <- probspace(rolldie(1),probs = c(1/12,3/12,1/12,3
   /12,1/12,3/12))
6
7 P_G1 <- fractions( S[4,"probs"] )
8 P_G2 <- fractions( S[5,"probs"] )
9 P_G3 <- fractions( S[6,"probs"] )
10 P_G4 <- P_G1+P_G2+P_G3
11 print(paste('The Probability is:',P_G4))
```

R code Exa 6.11.a Probability of students failed in maths test from a class of 120 students

```
1 #Page Number: 85
2 library(MASS)
3 Total <- 120
4 Maths101 <- 65
5 French101 <- 75
6 Both <- 35
7
8 mathpassprob <- fractions( Maths101 / Total )
9 mathfailprob <- 1 - mathpassprob
10 print(paste('Probability:',mathfailprob))
11
12 #The answer may vary due to difference in
   representation
```

R code Exa 6.11.b Probability of students passed in both test from a class of 120 students

```
1 #Page Number: 85
2
3 library(MASS)
4 Total <- 120
5 Maths101 <- 65
6 French101 <- 75
7 Both <- 35
8
9 bothpassprob <- fractions( Both / Total )
10 print(paste('Probability would be:',bothpassprob))
11
12 #The answer may vary due to difference in
    representation
```

R code Exa 6.11.c Probability of students failed in both test from a class of 120 students

```
1 #Page Number: 85
2
3 library(MASS)
4 Total <- 120
5 Maths101 <- 65
6 French101 <- 75
7 Both <- 35
8
9 mathpassprob <- fractions( Maths101 / Total )
10 frenchpassprob <- fractions( French101 / Total )
11 bothpassprob <- fractions( Both / Total )
12 allpassprob <- mathpassprob + frenchpassprob -
    bothpassprob
13 allfailprob <- 1 - allpassprob
14 print(paste('Probability :',allfailprob ))
```

```
15
16 #The answer may vary due to difference in
    representation
```

R code Exa 6.12 Probability of getting a card that is either red or a king from a deck of 52 cards

```
1 #Page Number: 86
2
3 library(prob)
4 library(MASS)
5
6 card <- cards()
7
8
9 diamond <- subset(card, suit == 'Diamond')
10 heart <- subset(card, suit == 'Heart')
11 E1 <- length(diamond$suit) + length(heart$suit)
12 probE1 <- fractions(E1 / length(card$suit))
13
14
15 E2 <- subset(card, rank == 'K')
16 probE2 <- fractions( length(E2$rank) / length(card$
    suit))
17
18
19 diaking <- intersect(diamond, E2)
20 heartking <- intersect(heart, E2)
21 intersec <- length(diaking$suit) + length(heartking$
    suit)
22 probintersec <- fractions(intersec / length(card$
    suit))
23
24 probunion <- probE1 + probE2 - probintersec
25 print(paste('The probability is:', probunion))
```

26 #The answer may vary due to difference in
representation.

R code Exa 6.13 Probability of getting atleast 1 head in tossing a fair coin
four times in a row

```
1 #Page Number: 86
2
3 library(prob)
4 library(MASS)
5
6 S <- iidspace(c("H","T"), ntrials = 4)
7
8 E <- S[16, 'probs']
9
10 probathead <- 1 - fractions(E)
11 print(paste('The probability is:',probathead))
```

R code Exa 6.14 Probability of drawing a card that is King and knowing
that there is no Queen in deck of 52 cards

```
1 #Page Number: 86
2 library(prob)
3 library(MASS)
4 card <- cards(makespace = T)
5
6 condprob <- fractions( Prob(card,rank == 'K' ,given
   = rank != 'Q') )
7 print(paste(" The Probability is:", condprob))
```

R code Exa 6.15 Probability of a student passed in maths test given that he or she is passed in atleast 1 test from a class of 120 students

```
1 #Page Number: 87
2
3 library(prob)
4 library(MASS)
5
6 Total <- 120
7 Maths101 <- 65
8 French101 <- 75
9 Both <- 35
10
11 A <- Maths101
12 P_A <- fractions( A / Total)
13
14 mathpassprob <- fractions( Maths101 / Total )
15 frenchpassprob <- fractions( French101 / Total )
16 bothpassprob <- fractions( Both / Total )
17 P_B <- mathpassprob + frenchpassprob - bothpassprob
18
19 condprob = P_A / P_B
20 print(paste("The probability is:", condprob))
21 #The answer may vary due to difference in
    representation.
```

R code Exa 6.16 Independent events of cards being drawn at random that are either red or a king from deck of 52 cards

```
1 #Page Number: 88
2
3 library(prob)
4 library(MASS)
5
6 card <- cards()
```

```

7
8 diamond <- subset(card, suit == 'Diamond')
9 heart <- subset(card, suit == 'Heart')
10
11
12 A <- length(diamond$suit) + length(heart$suit)
13 probA <- fractions(A / length(card$suit))
14
15 B <-subset(card, rank == 'K')
16 probB <- fractions( length(B$rank) / length(card$
    suit))
17
18 cat(paste("P(A)=" ,probA ,"\nP(B)=" ,probB))
19
20 diaking <- intersect(diamond, B)
21 heartking <-intersect(heart, B)
22 intersec <- length(diaking$suit) + length(heartking$
    suit)
23
24 probintersec <- fractions(intersec / length(card$
    suit))
25 cat(paste(probintersec))

```

R code Exa 6.17 Probability of a ship departing on time

```

1 #Page Number: 88
2
3 library(prob)
4
5 P_A <- 0.81
6
7 P_B <- 0.77
8
9 condAB <- 0.91
10

```

```

11 bayesBA <- (condAB * P_B) / P_A
12
13 print(paste('The probability is:', round(bayesBA, 3)))

```

R code Exa 6.18 Probability of getting a blue marble during first draw given that the second draw gave a blue marble from 2 boxes using bayes theorem

```

1 #Page Number: 89
2
3 library(prob)
4
5 S1 <- rep(c("Red", "Blue"), times = c(4,3))
6 Box1 <- urnsamples(S1, size = 1, replace = F,
7   ordered = F)
8 dfBox1 <- c(Box1)
9 totalred1 <- sum(dfBox1$out == "Red")
10 totalblue1 <- sum(dfBox1$out == "Blue")
11 prob_totalred1 <- totalred1 / length(dfBox1$out)
12 prob_totalblue1 <- totalblue1 / length(dfBox1$out)
13
14 S2 <- rep(c("Red", "Blue"), times = c(10,20))
15 Box2 <- urnsamples(S2, size = 1, replace = F,
16   ordered = F)
17 dfBox2 <- c(Box2)
18 totalred2 <- sum(dfBox2$out == "Red")
19 totalblue2 <- sum(dfBox2$out == "Blue")
20
21 f_red_totalred1 <- totalred1 - 1
22 f_red_totalred2 <- totalred2 + 1
23 red_totalmarble.in.1 <- length(dfBox1$out) - 1
24 red_totalmarble.in.2 <- length(dfBox2$out) + 1
25 f_prob_totalred1 <- f_red_totalred1 / red_
26   totalmarble.in.1
27 f_prob_totalred2 <- f_red_totalred2 / red_

```

```

    totalmarble.in.2
25 f_prob_totalblue1 <- totalblue1 / (length(dfBox1$out
    ) - 1)
26 f_prob_totalblue2 <- totalblue2 / (length(dfBox2$out
    ) + 1 )
27
28 s_blue_totalblue1 <- totalblue1 - 1
29 s_blue_totalblue2 <- totalblue2 + 1
30 blue_totalmarble.in.1 <- length(dfBox1$out) - 1
31 blue_totalmarble.in.2 <- length(dfBox2$out) + 1
32 s_prob_totalblue1 <- s_blue_totalblue1 / blue_
    totalmarble.in.1
33 s_prob_totalblue2 <- s_blue_totalblue2 / blue_
    totalmarble.in.2
34
35 bayes <- (s_prob_totalblue2 * prob_totalblue1 ) / ( (
    s_prob_totalblue2 * prob_totalblue1 ) + ( f_prob
    _totalblue2 * prob_totalred1 ))
36 print(paste("The probability is :",round(bayes,2)))

```

Chapter 7

Random Variables

R code Exa 7.1 Random variable operations on a rolling die

```
1 #Page Number: 93, 94(contd.), 95(contd.), 97(contd.)  
  , 99(contd.) .  
2  
3 library(prob)  
4 library(MASS)  
5 die <- rolldie(1,makespace = T)  
6 rx <- die$X1  
7 print(rx)  
8  
9 X_1 <- die[1,2]  
10 X_2 <- die[2,2]  
11 X_3 <- die[3,2]  
12 X_4 <- die[4,2]  
13 X_5 <- die[5,2]  
14 X_6 <- die[6,2]  
15  
16 print(paste('P(X = 1) =',fractions(X_1)))  
17 print(paste('P(X = 2) =',fractions(X_2)))  
18 print(paste('P(X = 3) =',fractions(X_3)))  
19 print(paste('P(X = 4) =',fractions(X_4)))  
20 print(paste('P(X = 5) =',fractions(X_5)))
```

```

21 print(paste('P(X = 6) =',fractions(X_6)))
22
23 E_X <- (die[1,1] * X_1 ) + ( die[2,1] * X_2 ) + (
      die[3,1] * X_3 ) + ( die[4,1] * X_4 ) + ( die
      [5,1] * X_5 ) + ( die[6,1] * X_6 )
24 print(E_X)
25
26 var <- ( ((die[1,1]-E_X)^2) * X_1 ) + ( ((die[2,1]-E
      _X)^2) * X_2 ) + ( ((die[3,1]-E_X)^2) * X_3 ) + (
      ((die[4,1]-E_X)^2) * X_4 ) + ( ((die[5,1]-E_X)
      ^2) * X_5 ) + ( ((die[6,1]-E_X)^2) * X_6 )
27 print(paste('Variance = ',round(var,4)))
28 sd <- sqrt(var)
29 print(paste('Standard Deviation = ',round(sd,4)))

```

R code Exa 7.2 Random variable operations on tossing a coin twice

```

1 #Page Number: 93, 94(contd.), 95(contd.), 97(contd.)
  , 99(contd.) .
2
3 library(prob)
4 library(MASS)
5
6 toss <- iidspace(c("H","T"), ntrials = 2)
7
8 X_0 <- toss[4,'probs']
9 X_1 <- toss[2,'probs'] + toss[3,'probs']
10 X_2 <- toss[1,'probs']
11 print(paste('P(X = 0) =',fractions(X_0)))
12 print(paste('P(X = 1) =',fractions(X_1)))
13 print(paste('P(X = 2) =',fractions(X_2)))
14
15 E_X <- (0 * X_0) + (1 * X_1) + (2 * X_2)
16 print(E_X)
17

```

```

18 var <- (((0 - E_X)^2)*(X_0)) + (((1 - E_X)^2)*(X_1))
      + (((2 - E_X)^2)*(X_2))
19 print(paste('Variance = ',var))
20 sd <- sqrt(var)
21 print(paste('Standard Deviation ',round(sd,4)))

```

R code Exa 7.3 Random variable operations in tossing a fair coin twice and then rolling a balanced die

```

1 #Page Number: 94, 95(contd.), 97(contd.), 99(contd.)
2
3 library(MASS)
4
5 S <- data.frame(TOSS = c('H','H','H','H','H','H','T',
      'T','T','T','T','T'),DIE = c
      (1,2,3,4,5,6,1,2,3,4,5,6))
6
7 X_0 <- sum(S$TOSS == "T") / length(S$TOSS)
8 X_1 <- sum(S$TOSS == "H") / length(S$TOSS)
9 print(paste('P(X = 0) =',fractions(X_0)))
10 print(paste('P(X = 1) =',fractions(X_1)))
11
12 Y_1 <- sum(S$DIE == "1") / length(S$DIE)
13 Y_2 <- sum(S$DIE == "2") / length(S$DIE)
14 Y_3 <- sum(S$DIE == "3") / length(S$DIE)
15 Y_4 <- sum(S$DIE == "4") / length(S$DIE)
16 Y_5 <- sum(S$DIE == "5") / length(S$DIE)
17 Y_6 <- sum(S$DIE == "6") / length(S$DIE)
18
19 mu_X <- (0 * X_0) + (1 * X_1)
20 print(mu_X)
21 mu_Y <- (1 * Y_1) + (2 * Y_2) + (3 * Y_3) + (4 * Y_
      4) + (5 * Y_5) + (6 * Y_6)
22 print(mu_Y)

```

```

23
24 varX <- (((0- mu_X)^2) * (X_0)) + (((1 - mu_X)^2) *
      (X_1))
25 VarY <- (((1 - mu_Y)^2) * (Y_1)) + (((2 - mu_Y)^2) *
      (Y_2)) + (((3 - mu_Y)^2) * (Y_3)) + (((4 - mu_Y)
      ^2) * (Y_4)) +
26       (((5 - mu_Y)^2) * (Y_5)) + (((6 - mu_Y)^2) *
      (Y_6))
27 print(paste('Variance:',varX, ' & ',round(VarY,4)))
28
29 sdX <- sqrt(varX)
30 sdY <- sqrt (VarY)
31 print(paste('Standard Deviation:',sdX, '&',round(sdY
      ,4)))
32
33 #The answer provided in the textbook is wrong

```

R code Exa 7.4 Random variable operations on a sample of 3 voters

```

1 #Page Number: 94, 95(contd.), 97(contd.), 99(contd.)
2
3 library(prob)
4 library(MASS)
5
6 vote <- iidspace(c("Y","N"), ntrials = 3)
7 print(vote)
8
9 X_0 <- vote[8,"probs"]
10 X_1 <- vote[4,"probs"] + vote[6,"probs"] + vote[7,"
      probs"]
11 X_2 <- vote[2,"probs"] + vote[3,"probs"] + vote[5,"
      probs"]
12 X_3 <- vote[1,"probs"]
13

```

```

14 print(paste("P(X = 0) =", fractions(X_0)))
15 print(paste("P(X = 1) =", fractions(X_1)))
16 print(paste("P(X = 2) =", fractions(X_2)))
17 print(paste("P(X = 3) =", fractions(X_3)))
18
19 E_X <- (0 * X_0) + (1 * X_1) + (2 * X_2) + (3 * X_3)
20 print(E_X)
21
22 var <- (((0 - E_X)^2)*(X_0)) + (((1 - E_X)^2)*(X_1))
      + (((2 - E_X)^2)*(X_2)) + (((3 - E_X)^2)*(X_3))
23 print(paste('Variance = ', var))
24 sd <- sqrt(var)
25 print(paste('Standard Deviation ', round(sd, 2)))

```

R code Exa 7.6 To find all possible grade pairs for a certain mark obtained

```

1 #Page Number: 95
2
3 library(tidyverse)
4 grade_combo <- function(x) {
5   dataframe <- data.frame(g1 = c("A", "B", "C", "D",
6     "F"),
7     g2 = c("A", "B", "C", "D", "F"))
8   %>%
9   expand.grid(.) %>%
10  data.frame(.) %>%
11  mutate(p = case_when(g1 == "A" ~ 4,
12    g1 == "B" ~ 3,
13    g1 == "C" ~ 2,
14    g1 == "D" ~ 1,
15    TRUE ~ 0),
16    p2 = case_when(g2 == "A" ~ 4,
17    g2 == "B" ~ 3,
18    g2 == "C" ~ 2,
19    g2 == "D" ~ 1,

```

```
18                                     TRUE ~ 0),
19         GPA = (p + p2)/2)
20
21
22     res <- dataframe %>%
23       filter(GPA == x)
24
25     return(res = res)
26 }
27
28 grade <- grade_combo(2.5)
29
30 print(grade[c(1,2)])
```

Chapter 8

Binomial Experiments

R code Exa 8.1 Probability of answering correctly in a pop quiz

```
1 #Page Number: 103, 106(contd.) .
2
3 library(prob)
4 library(MASS)
5 see <- iidspace(c("C","I"), ntrials = 7)
6 options(max.print = 10000)
7 print(see)
8
9 size <- 7
10 prob <- 1/2
11
12 X_0 <- dbinom(0,size,prob)
13 X_1 <- dbinom(1,size,prob)
14 X_2 <- dbinom(2,size,prob)
15 X_3 <- dbinom(3,size,prob)
16 X_4 <- dbinom(4,size,prob)
17 X_5 <- dbinom(5,size,prob)
18 X_6 <- dbinom(6,size,prob)
19 X_7 <- dbinom(7,size,prob)
20
21 print(paste('P(X = 0) =',round(X_0,3)))
```

```
22 print(paste('P(X = 1) =', round(X_1, 3)))
23 print(paste('P(X = 2) =', round(X_2, 3)))
24 print(paste('P(X = 3) =', round(X_3, 3)))
25 print(paste('P(X = 4) =', round(X_4, 3)))
26 print(paste('P(X = 5) =', round(X_5, 3)))
27 print(paste('P(X = 6) =', round(X_6, 3)))
28 print(paste('P(X = 7) =', round(X_7, 3)))
```

R code Exa 8.2 Probability of finding mortar shell explosion on enemy target

```
1 #Page Number: 104, 107(contd.) .
2
3 fprob <- 0.10
4 size <- 20
5 sprob <- 1 - fprob
6
7 Xles14 <- dbinom(0:14, size = size, prob = sprob)
8 probXles14 <- sum(Xles14)
9 more15 <- 1 - probXles14
10 print(round(more15, 3))
```

R code Exa 8.3 Probability of at least 1 sensor detecting gas leak

```
1 #Page Number: 104, 107(contd.) .
2
3 sprob <- 0.80
4 size <- 5
5 X_0 <- dbinom(0, size, sprob)
6 res <- 1 - X_0
7 print(round(res, 1))
```

R code Exa 8.4 Probability of fire accident due to arson

```
1 #Page Number: 104, 108(contd.) .
2
3 sprob <- 0.25
4 size <- 10
5
6
7 less4 <- dbinom(0:4, size, sprob )
8 res <- sum(less4)
9 print(round(res,3))
```

R code Exa 8.5 Probability of members who are seeking capital punishment

```
1 #Page Number: 104, 108(contd.) .
2
3 probagainst <- 0.70
4 size <- 12
5 probfavour <- 1- probagainst
6
7 less6 <- dbinom(0:6, size, probfavour)
8 probless6 <- sum(less6)
9 more7 <- 1 - probless6
10 print(round(more7,3))
```

R code Exa 8.6 Tossing a fair coin 15 times

```
1 #Page Number: 104
```

```
2
3 library(prob)
4
5 a <- tosscoin(15)
6 n <- ncol(a)
7 print(n)
8
9 print(levels(a$toss1))
10
11 b <- tosscoin(1)
12 c <- length(b[b$toss1 == 'H']) / length(b$toss1)
13 print(c)
```

Chapter 9

Normal Curve and Normal Distribution

R code Exa 9.1 Percentage of children facing difficulties with new teaching method

```
1 #Page Number: 119, 123 (contd.) .  
2  
3 IQlev <- 95  
4 prevm <- 112  
5 prevsd <- 7.8  
6  
7 res<- pnorm(IQlev, mean = prevm, sd = prevsd)  
8 print(round(res,4))
```

R code Exa 9.2.a Probability for a standard normal random variable

```
1 #Page Number: 122  
2  
3  
4 a <- 1.5
```

```
5
6 res <- 0.5 - pnorm(a, lower.tail = F)
7 print(round(res,4))
```

R code Exa 9.2.b Probability for a standard normal random variable

```
1 #Page Number: 122
2
3 a <- -2.33
4
5 res <- 1 - (2 *pnorm(a, lower.tail = TRUE ))
6 print(round(res,4))
```

R code Exa 9.2.c Probability for a standard normal random variable

```
1 #Page Number: 122
2
3 a <- 1.69
4
5 res <- 1 - pnorm(a, lower.tail = FALSE)
6 print(round(res,4))
```

R code Exa 9.2.d Probability for a standard normal random variable

```
1 #Page Number: 122.
2
3 a <- -1.25
4
5 res <- 1 - pnorm(a)
6 print(round(res,4))
```

R code Exa 9.3 Percentage of day time temperature between 80 to 100

```
1 #Page Number: 123
2
3 mean <- 89
4 sd <- 6.9
5 x1 <- 80
6 x2 <- 100
7
8 a <- (x1 - mean)/sd
9 b <- (x2 - mean)/sd
10
11 res <- pnorm(round(b,2)) - pnorm(round(a,2))
12 print(round(res,4))
```

Chapter 10

Applications of the Normal Distribution

R code Exa 10.1 Probability of 6 or more people suffering from hypothermia

```
1 #Page Number: 131
2
3 prob <- 0.75
4 n <- 8
5
6 res <- dbinom(6:8,n,prob)
7 print(sum(round(res,3)))
```

R code Exa 10.2 Probability of 80 or less people suffering from hypothermia

```
1 #Page Number: 132, 133(contd.)
2
3
4 prob <- 0.75
```

```

5 n <- 120
6
7
8 mu <- n*prob
9 sigma <- round(sqrt((n*prob)*(1-prob)),4)
10
11 res <- round(pnorm(80 + 0.5, mean = mu, sd = sigma)
12           ,4)
13 print(res)

```

R code Exa 10.3.a Probability of 640 or more students passing the exam

```

1 #Page Number: 134
2
3 prob <- 0.57
4 n <- 950
5
6 mu <- n*prob
7 sigma <- round(sqrt((n*prob)*(1-prob)),4)
8
9 res <- pnorm(640 - 0.5, mean = mu, sd = sigma)
10 print(floor(res))

```

R code Exa 10.3.b Probability of 600 or fewer students passing the exam

```

1 #Page Number: 134
2
3 prob <- 0.57
4 n <- 950
5
6 mu <- n*prob
7 sigma <- round(sqrt((n*prob)*(1-prob)),4)
8

```

```
9 res <- pnorm(600 + 0.5, mean = mu, sd = sigma)
10 print(round(res,1))
```

R code Exa 10.3.c Probability of students passing the exam between 585 and 625

```
1 #Page Number: 134
2
3 prob <- 0.57
4 n <- 950
5
6 mu <- n*prob
7 sigma <- sqrt((n*prob)*(1-prob))
8
9 X_grt585 <- ((585 - 0.5) - mu) / sigma
10 X_les625 <- ((625 + 0.5) - mu) /sigma
11
12 res<- pnorm(X_grt585)
13 print(round(res,1))
```

R code Exa 10.4 Probability of sample batteries having average life of 52 or less

```
1 #Page Number: 137
2
3 mu_x <- 54
4 n <- 50
5 sigma <- 6
6 sigma_x <- round(sigma / sqrt(n),2)
7
8
9 res<- pnorm(52,mean = mu_x, sd = sigma_x)
10 print(round(res,4))
```

R code Exa 10.5.a Distribution of sample average of bacteria count per ML of milk

```
1 #Page Number: 137
2
3 mu_x <- 2550
4 n <- 42
5 sigma <- 300
6
7 sigma_x <- round(sigma / sqrt(n),2)
8
9 print(sigma_x)
```

R code Exa 10.5.b Probability of sample having bacteria between 2400 to 2700

```
1 #Page Number: 137
2
3 mu_x <- 2550
4 n <- 42
5 sigma <- 300
6 sigma_x <- round(sigma / sqrt(n),2)
7
8
9 xless <- ((2400 - mu_x) / sigma_x)
10 xgrtr <- ((2700 - mu_x) / sigma_x)
11
12 res <- pnorm(xgrtr) - pnorm(xless)
13 print(round(res,4))
```

R code Exa 10.6 Probability of sample average of IQ scores not deffering from population mean by more than 1 point

```
1 #Page Number: 138
2
3 mu_x <- 104
4 n <- 64
5 sigma <- 5
6 sigma_x <- sigma / sqrt(n)
7
8 xless <- ((103 - mu_x) / sigma_x)
9 xgrtr <- ((105 - mu_x) / sigma_x)
10
11 res <- pnorm(xgrtr) - pnorm(xless)
12 print(round(res,4))
```

Chapter 11

Estimation of Population Parameters

R code Exa 11.2 Computing a 90 percent confidence interval for mean PE ratio of all utility stocks in NYSE

```
1 #Page Number: 151
2
3 data <- data.frame( "Utility Comp." = c(1,2,3,4,5) ,
4   "P/E Ratio" = c(20.41,12.69,16.15,10.09,13.10) )
5
6 conf.interval = function(data, conf) {
7   n <- length(data)
8   alpha <- 1 - conf
9   E <- qt((alpha/2) + conf, df = n-1)
10  xbar <- mean(data)
11  sdx <- sqrt(var(data)/length(data))
12  c(xbar - E * sdx, xbar + E * sdx)
13 }
14 print(round(conf.interval(data$P.E.Ratio, 0.90),2))
```

R code Exa 11.3 Computing 95 percent confidence interval for the mean January average temperature in Phoenix

```
1 #Page Number: 152.
2
3 data <- read.csv("Ex11_3.csv")
4
5 conf.interval <- function(data, conf) {
6   n <- length(data)
7   alpha <- 1 - conf
8   E <- qnorm((alpha/2) + conf)
9   xbar <- mean(data)
10  sdx <- sqrt(var(data)/length(data))
11  c(xbar - E * sdx, xbar + E * sdx)
12 }
13 res <- conf.interval(data$Average.temperature, 0.95)
14 print(round(res,2))
```

R code Exa 11.4 Computing 90 percent confidence interval for the percentage of US parents whose kids favourite food is pizza

```
1 #Page Number: 159
2
3 n <- 1034
4 X <- 848
5
6 conf.interval <- function(data, conf, x){
7   p <- x/data
8   alpha <- 1 - conf
9   SE <- sqrt(p*(1 - p)/data)
10  E <- qnorm((alpha / 2) + conf)*SE
11  c(p - E, p + E )
12 }
13 res <- conf.interval(n,0.90,X)
14 print(round(res,4))
```


Chapter 12

Hypothesis Testing for a Single Population

R code Exa 12.2 Hypothesis testing on claim of mean household income in a business district

```
1 #Page Number: 176, 190(contd.) .
2
3
4 tstatvalue <-function(n,x,p){
5 pbar <- round(x/n,4)
6 del <- (sqrt(n)*(pbar -p)) / sqrt(p*(1-p))
7 del
8 }
9 tstat <-round(tstatvalue(72,15,0.25),4)
10 print(paste("Test Statistic:",tstat))
11
12 res <- round(pnorm(-abs(tstat)),4)
13 print(paste("P-value :",res))
```

R code Exa 12.3 Hypothesis testing on claims of nutritionist on the rise of potato consumption

```

1 #Page Number: 179, 190(contd.).
2
3 n <- 46
4 xbar <- 53.4
5 mu <- 48.3
6 sd <- 4.12
7
8 alpha = .05
9 t.alpha = round(qt(1-alpha, df=n-1),4)
10
11 tstat <- round((sqrt(n)*(xbar - mu)) / sd,4)
12 print(paste("The test statistic",tstat))
13
14 res <- round(pnorm(-abs(tstat)),4)
15 print(paste("P-value :",res))

```

R code Exa 12.4 Hypothesis testing on claims based on students councils on time spent in library

```

1 #Page Number: 183, 190(contd.).
2
3 hour <- c(10.4,9.1,13.6,6.4,7.9,8.6,11.8,12.6)
4 mu <- 12.5
5 xbar <- mean(hour)
6 s <- round(sd(hour),4)
7 del <- round((sqrt(length(hour))*(xbar-mu)) / s,4)
8
9 alpha5 <- 0.05
10 t.alpha5 <- round(qt(alpha5/2, df = length(hour) -1,
    lower.tail = F),3)
11
12 alpha1 <- 0.01
13 t.alpha1 <- round(qt(alpha1/2, df = length(hour) -1,
    lower.tail = F),3)
14

```

```

15 cat("Test Statistics:", abs(del))
16
17 n = length(hour)
18 p = pnorm(-abs(del))
19 print(paste("P-value:", round(p,3)))
20
21 #The answer provided in the textbook is wrong.

```

R code Exa 12.5 Hypothesis testing on claims of management on share of fruits is 50 percent

```

1 #Page Number: 185, 191(contd.).
2
3 data <- read.csv("EX12_5.csv")
4 n <- length(data$Diameters)
5 p <- 0.50
6 large <- data$Diameters > 3.0
7 pbar <- sum(large)/ n
8 del <- (sqrt(n)*(pbar -p)) / sqrt(p*(1-p))
9
10 zt <- round(qnorm(0.05),3)
11 print(paste("The test statistic",round(del,4)))
12
13 res <- round(pnorm(-abs(del)),4)
14 print(paste("P-value :",res))
15 print(paste("Combined Area , P-Value:",res+res))

```

R code Exa 12.6 Hypothesis testing on claims on weight of the newly minted coins

```

1 #Page Number: 187, 192(contd.).
2
3

```



```

4 data <- data.frame("Weight" = c
  (2.99,3.01,3.03,3.05,3.07,3.09,3.11,3.13,3.15,3.17,3.19)
  , "Frequency" = c(1,4,4,4,7,17,24,17,13,6,2))
5 n <- 100
6 sig <- 0.05
7 interval <- seq( from = 2.98, to = 3.22, by = 0.02)
8 subint <- cut(data$Weight,interval)
9 redata <- data.frame("Mid" = c(2.99,3.01,3.03,3.05,
  3.07, 3.09, 3.11, 3.13, 3.15, 3.17, 3.19,3.21), "
  Frequency" = c(1,4,4,4,7,17,24,17,13,6,2,1) )
10 rf <- redata$Frequency/ sum(redata$Frequency)
11 xbar <- sum(redata$Mid * rf)
12 s <- sum(((redata$Mid-xbar)^2)*redata$Frequency)
13 s2 <- s / (sum(redata$Frequency)-1)
14
15
16 del <- round(((n-1)*s2) / (sig)^2,4)
17 chi <- round(qchisq(0.10, df = n-1, lower.tail = F)
  ,4)
18
19 print(paste("The test statistic:",del))
20
21 res <- pchisq(del,df = n-1,lower.tail = F)
22 print(paste("P-value :",round(res,3)))

```

R code Exa 12.7 Verifying Normality for the given dataset

```

1 #Page Number: 194
2
3 hour <- c(10.4,9.1,13.6,6.4,7.9,8.6,11.8,12.6)
4 xbar <- mean(hour)
5 s <-round(sd(hour),2)
6 Z.i <- sort(round(as.vector((hour - xbar)/ s),2))
7 B.i <- round(sort(pnorm(Z.i)),4)
8 A.i <- round(1:length(hour) / length(hour),4)

```

```
9 A.i.min1 <- round(0:(length(hour)-1) / length(hour)
,4)
10 C.i <- abs(round(A.i - B.i,4))
11 C.i.str <- abs(round(A.i.min1 - B.i,4))
12
13 dt <- data.frame("Zi" = Z.i, "Bi" = B.i, "Ai" = A.i,
" Ai_minus_1" = A.i.min1, "Ci" = C.i, "Ci_str" =
C.i.str)
14 print(dt)
15
16 del_ks <- round(max(C.i),4)
17 print(del_ks)
```

Chapter 13

Hypothesis Testing to Compare Two Populations

R code Exa 13.2 Testing the claims on painkiller which are identical in terms of mean time required for body absorption

```
1 #Page Number: 212, 224(contd.) .
2
3 n1 <- 10
4 n2 <- 10
5 xbar1 <- 20.2
6 xbar2 <- 17.9
7 s1 <- 8.1
8 s2 <- 7.3
9
10 poolsample <- round( sqrt(((n1-1)*(s1)^2 + (n2-1)*(
    s2)^2 ) / (n1+n2-2) ),4)
11 cat("Pooled Sample :",poolsample)
12
13 tstat <- round((xbar1-xbar2) /( poolsample* sqrt((1/
    n1) + (1/n2))),4)
14 cat("Test Statistic value:",tstat)
15
16 alpha <- 0.05
```

```

17 t.alpha <- round(qt(alpha/2, df = n1+n2-2, lower.tail = F),3)
18 cat(tstat,"<",t.alpha)
19
20 res <- 2 * pt(-abs(tstat),df= n1+n2-2)
21 cat("The P-Value:",round(res,2))

```

R code Exa 13.3 Testing the difference between the mean stopping distances of two types of tyres

```

1 #Page Number: 213, 224(contd.).
2
3 n1 <- 10
4 n2 <- 12
5 xbar1 <- 51
6 xbar2 <- 55
7 s1 <- 8
8 s2 <- 3
9
10 tstat <- abs(round((xbar1 - xbar2) / sqrt(((s1)^2/n1
    ) + ((s2)^2/n2)),3))
11 q <-round( (((s1)^2/n1)+((s2)^2/n2))^2 / ((s1)^4 /
    ((n1)^2*(n1-1)) + (s2)^4 / ((n2)^2*(n2-1))),0)
12 cat("Test Statistic value:",tstat)
13 cat("q:",q)
14
15 alpha <- 0.05
16 t.alpha <- round(qt(alpha/2, df = q,lower.tail = F)
    ,3)
17 cat(tstat,"<",t.alpha)
18
19 res <- 2 * pt(-abs(tstat),df= q)
20 cat("The P-Value:",round(res,2))

```

R code Exa 13.4 Testing on effects of a new hypertension drug which changes the body temperature

```
1 #Page Number: 215, 225(contd.).
2
3 n <- 7
4 bdrug <- c(98.9,100.2,97.9,99.4,98.1,98.5,101.8)
5 adrug <- c(98.6,99.7,97.6,99.9,98.2,97.9,99.2)
6 diff <- bdrug - adrug
7 dbar <- round(sum(diff) / n,4)
8 sD <- sqrt (sum((diff - dbar)^2) / (n-1))
9
10 tstat <- round((sqrt(n)*dbar) / (sD),4)
11 cat("Test Statistic value:",tstat)
12
13 alpha <- 0.05
14 t.alpha <- round(qt(alpha/2,df = n-1,lower.tail = F)
15 ,3)
16 cat(tstat,"<",t.alpha)
17
18 res <- 2 * pt(-abs(tstat),df= n-1)
19 cat("The P-Value:",round(res,2))
```

R code Exa 13.5 Testing the beliefs on the methods which is superior than other in treating patients with broken arm

```
1 #Page Number: 218, 225(contd.).
2
3 n1 <- 145
4 n2 <- 111
5 x1 <- 21
6 x2 <- 9
```

```

7 pt.est1 <- round(x1/n1,4)
8 pt.est2 <- round(x2/n2,4)
9 pcap <- round((x1+x2) / (n1+n2),4)
10
11 tstat <- round((pt.est1 - pt.est2) / sqrt( pcap*(1-
      pcap)*((1/n1)+(1/n2)) ),4)
12 cat("Test Statistic value:",tstat)
13
14 zt <- round(qnorm(0.05,lower.tail = F),3)
15 cat(tstat,"<",zt)
16
17 res <- pnorm(-tstat)
18 cat("The P-Value:",round(res,4))

```

R code Exa 13.6 Testing on data on ball bearing from 2 machines indicating that the machines are identical

```

1 #Page Number: 220, 225(contd).
2
3 n1 <- 10
4 n2 <- 10
5 xbar1 <- 1.96
6 xbar2 <- 2.01
7 s1 <- 0.031
8 s2 <- 0.050
9
10 k1 <- n1-1
11 k2 <- n2-1
12 tstat_F <- (s1)^2 / (s2)^2
13 cat("Test Statistic value:",tstat_F)
14
15 alpha <- 0.05
16 f.test_r <- round( qf(alpha/2,df1 = k1,df2 = k2,
      lower.tail = F),2)
17 f.test_l <- round(qf(alpha/2,df1 = k1,df2 = k2),3)

```

```
18 cat(f.test_l,"<",tstat_F,"<",f.test_r)
19
20 right_side <- round(1/tstat_F,3)
21 res = 2 * round(pf(1/right_side,df1 = k1,df2 = k2,
    lower.tail = T),1)
22
23 cat("The P-Value:",res)
```

Chapter 14

Bivariate Quantitative Data Correlation and Regression

R code Exa 14.1 Computing the operations of correlation and regression on population consisting of all married couples

```
1 #Page Number: 241, 246(contd.), 251(contd.), 255(
  contd.), 257(contd.).
2
3 data <- data.frame("Couple No." = c(1:12), "Husband
  Age" = c(30,29,36,72,37,36,51,48,37,50,51,36),"
  Wife Age" = c
    (27,20,34,67,35,37,50,46,36,42,46,35))
4 n <- length(data$Couple.No.)
5
6 hus_interval <- seq( from = 28.5, to = 72.5, by =
  11)
7 hus_class <- cut(data$Husband.Age, hus_interval)
8 hus_frequency <- cbind(table(hus_class))
9 hus.Rel_Freq <- (hus_frequency/n)*100
10 df1 <- data.frame("Group" = unique(hus_class), "
  Frequency" = hus_frequency, "Relative.Frequency"
  = round(hus.Rel_Freq,2))
11 print(df1)
```



```

12
13 wif_interval <- seq( from = 19.5, to = 67.5, by =
    12)
14 wif_class <- cut(data$Wife.Age, wif_interval)
15 wif_frequency <- cbind(table(wif_class))
16 wif.Rel_Freq <- (wif_frequency/n)*100
17 df2 <- data.frame("Group" = unique(wif_class), "
    Frequency" = wif_frequency, "Relative.Frequency"
    = round(wif.Rel_Freq,2))
18 print(df2)
19
20
21
22 xbar <- sum(data$Husband.Age) / n
23 ybar <- round(sum(data$Wife.Age) / n,2)
24 S_x <- round(sd(data$Husband.Age),4)
25 S_y <- round(sd(data$Wife.Age),4)
26 sample.average <- data.frame( "Couple" = c("Husband"
    ,"Wife"), "Mean" = c(xbar,ybar), "SD" = c(S_x,S_y
    ))
27 print(sample.average)
28
29 plot(data$Husband.Age,data$Wife.Age)
30 cor_coef <- round(cor(data$Husband.Age,data$Wife.Age
    ),4)
31 cat("The Correlation Coefficient is",cor_coef)
32
33 tstat.r <- round((cor_coef*(sqrt(n-2))) / sqrt(1-(
    cor_coef)^2),4)
34 cat("Test Statistic value:",tstat.r)
35
36 alpha <- 0.05
37 critical.value <-round(qt(alpha,df = n-2,lower.tail
    = F),3)
38 cat(tstat.r,">",critical.value)
39
40 Z <- round(qnorm(alpha/2,lower.tail = F),2)
41 C <- round(exp((2/sqrt(n-3))*(Z)),3)

```

```

42 D <- round((1+cor_coef)/(1-cor_coef),4)
43 A <- C*D
44 B <- D/C
45 lower <- round((B-1)/(B+1),4)
46 upper <- round((A-1)/(A+1),4)
47 cat("95% confidence interval is",lower,"",upper)
48
49 reg.fun <- function(data1,data2){
50   beta1 <- round(cor(data1,data2)*(sd(data2)/sd(
      data1)),4)
51   beta0 <- round(mean(data2)-(beta1*mean(data1)),3)
52   Y.fit <- round(beta0 + beta1*(data1),3)
53   return(data2 - Y.fit)
54 }
55 sumation.of.Y_fit <- reg.fun(data$Husband.Age,data$
      Wife.Age)
56 Std_error <- round(sqrt(sum((sumation.of.Y_fit)^2) /
      (n-2)),4)
57 print(paste("Standard Error:",Std_error))
58
59 x <- 62
60 beta1 <- round(cor_coef*(S_y/S_x),4)
61 beta0 <- round(ybar-(beta1*xbar),3)
62 yfit <- beta0 + beta1*(62)
63 cat("Husband's Age is 62 years ,prediction for wife's
      age:",yfit)
64
65 E <- round(critical.value * Std_error * sqrt(1 + (1/
      n)+((x-xbar)^2)/((n-1) * (S_x)^2)),2)
66 cat("Error Margin:",E)
67 cat("90% prediction:",yfit-E,"",yfit+E )

```

R code Exa 14.2 Computing the operations of correlation and regression on population of sales of hot chocolate and temperature for a snack shop

```

1 #Page Number: 244, 251(contd.), 256(contd.), 257(
  contd.).
2
3 data <- data.frame("Match No." = c(1:9),"Daytime
  Temp." = c(51,60,65,71,39,32,81,76,66),"Hot
  Chocolate Sale"= c
  (187,210,137,136,241,262,110,143,152))
4 n <- length(data$Match.No.)
5 plot(x = data$Daytime.Temp., y = data$Hot.Chocolate.
  Sale, xlim = c(0,90), ylim = c(0,270))
6
7 cor_coef <- round(cor(data$Daytime.Temp.,data$Hot.
  Chocolate.Sale),4)
8 cat("The Correlation Coefficient is",cor_coef)
9
10 tstat.r <- round(((cor_coef*(sqrt(n-2))) / sqrt(1-(
  cor_coef)^2),4)
11 cat("Test Statistic value:",tstat.r)
12 alpha <- 0.05
13 critical.value <-round(qt(alpha,df = n-2,),3)
14 cat(tstat.r,"<",critical.value)
15
16 xbar <- mean(data$Daytime.Temp.)
17 ybar <- mean(data$Hot.Chocolate.Sale)
18 s_x <- sd(data$Daytime.Temp.)
19 s_y <- sd(data$Hot.Chocolate.Sale)
20 beta1 <- round((cor_coef * (s_y/s_x)),4)
21 beta0 <- round(ybar-(beta1*xbar),4)
22 cat("Regression line:",beta0,"-(",-beta1,")X")

```

Chapter 15

Bivariate Categorical Data Contingency Table

R code Exa 15.1 Testing independence of gender wise usage of 2 types of painkillers

```
1 #Page Number: 274
2
3 data <- matrix(c(115,85,135,165),nrow = 2,ncol = 2,
  byrow = T)
4 data <- rbind(data, data[1,]+data[2,])
5 data <- cbind(data, data[,1]+data[,2])
6 rownames(data) <- c("Male", "Female", "Column Total")
7 colnames(data) <- c("Med-1", "Med-2", "Row Total")
8 n <- data[3,3]
9 data
10
11 E11 <- data[1,3]*data[3,1] / n
12 E12 <- data[1,3] - E11
13 E21 <- data[3,2] - E11
14 E22 <- data[2,3] - E21
15
16 tstat.c <- ((data[1,1] - E11)^2 / E11) + ((data[1,2]
  - E12)^2 / E12) + ((data[2,1] - E21)^2 / E21) +
```

```

      ((data[2,2] - E22)^2 / E22)
17 cat("test statistic:",tstat.c)
18
19 alpha <- 0.05
20 a <- 2
21 b <- 2
22 k <- (a-1)*(b-1)
23
24 critical.val <- round(qchisq(alpha,k,lower.tail = F)
      ,3)
25 cat(tstat.c,">",critical.val)

```

R code Exa 15.2 Testing independence on whether smoking causes hypertension on dataset of randomly selected set of patients

```

1 #Page Number: 278
2
3 data <- matrix(c(20,38,28,50,27,18),nrow = 2,ncol =
      3,byrow = T)
4 data <- rbind(data, data[1,]+data[2,])
5 data <- cbind(data, data[,1]+data[,2]+data[,3])
6 rownames(data) <- c("Hypertension", "No hypertension",
      ", "Column Total")
7 colnames(data) <- c("Non Smoker", "Moderate", "Heavy",
      "Row Total")
8 n <- data[3,4]
9 print(data)
10
11 E11 <- round(data[1,4]*data[3,1] / n,2)
12 E12 <- round(data[1,4]*data[3,2] / n,2)
13 E13 <- round(data[1,4]*data[3,3] / n,2)
14 E21 <- round(data[2,4]*data[3,1] / n,2)
15 E22 <- round(data[2,4]*data[3,2] / n,2)
16 E23 <- round(data[2,4]*data[3,3] / n,2)
17

```

```

18 tstat.c <- round(((data[1,1] - E11)^2 / E11) + ((
    data[1,2] - E12)^2 / E12)+((data[1,3] - E13)^2 /
    E13) + ((data[2,1] - E21)^2 / E21) + ((data[2,2]
    - E22)^2 / E22) +((data[2,3] - E23)^2 / E23),4)
19 cat("test statistic:",tstat.c)
20
21 alpha <- 0.05
22 a <- 2
23 b <- 3
24 k <- (a-1)*(b-1)
25
26 critical.val <- round(qchisq(alpha,k,lower.tail = F)
    ,3)
27 print(critical.val)

```

R code Exa 15.3 Testing independence of accidents involved in accordance with BAC levels and the region

```

1 #Page Number: 280
2
3 data <- matrix(c
    (5281,525,3064,3693,336,1949,5337,519,3213,10967,979,6722)
    ,nrow = 4,ncol = 3,byrow = T)
4 data <- rbind(data, data[1,]+data[2,]+data[3,]+data
    [4,])
5 data <- cbind(data, data[,1]+data[,2]+data[,3])
6 rownames(data) <- c("West", "North-East", "Mid-West", "
    South", "Column Total")
7 colnames(data) <- c("No Alcohol", "Low Alcohol", "High
    Alcohol", "Row Total")
8 n <- data[5,4]
9 data
10
11 E11 <- round(data[1,4]*data[5,1] / n,2)
12 E12 <- round(data[1,4]*data[5,2] / n,2)

```

```

13 E13 <- round(data[1,4]*data[5,3] / n,2)
14
15 E21 <- round(data[2,4]*data[5,1] / n,2)
16 E22 <- round(data[2,4]*data[5,2] / n,2)
17 E23 <- round(data[2,4]*data[5,3] / n,2)
18
19 E31 <- round(data[3,4]*data[5,1] / n,2)
20 E32 <- round(data[3,4]*data[5,2] / n,2)
21 E33 <- round(data[3,4]*data[5,3] / n,2)
22
23 E41 <- round(data[4,4]*data[5,1] / n,2)
24 E42 <- round(data[4,4]*data[5,2] / n,2)
25 E43 <- round(data[4,4]*data[5,3] / n,2)
26
27 tstat.c <- round((((data[1,1] - E11)^2 / E11) + ((
      data[1,2] - E12)^2 / E12)+((data[1,3] - E13)^2 /
      E13) + ((data[2,1] - E21)^2 / E21) + ((data[2,2]
      - E22)^2 / E22) +((data[2,3] - E23)^2 / E23) +((
      data[3,1] - E31)^2 / E31) + ((data[3,2] - E32)^2
      / E32)+((data[3,3] - E33)^2 / E33)+((data[4,1] -
      E41)^2 / E41) + ((data[4,2] - E42)^2 / E42)+((
      data[4,3] - E43)^2 / E43),4)
28 cat("test statistic:",tstat.c)
29
30 alpha <- 0.05
31 a <- 4
32 b <- 3
33 k <- (a-1)*(b-1)
34 critical.val <- round(qchisq(alpha,k,lower.tail = F)
      ,3)
35 cat(tstat.c,">",critical.val)

```

Chapter 16

Multinomial Experiments Goodness of Fit Test

R code Exa 16.1 Correlation coefficient describing the probability distribution of the outcome when 15 marbles are drawn at random from a box of 100 where 60 is red and 40 is blue

```
1 #Page Number: 289
2
3 library(ggplot2)
4
5 x <- 0:15
6 y <- 15 - x
7 xpy <- data.frame(X = x, Y = y)
8 coords = paste(x,y,sep=" ")
9
10 ggplot(xpy, aes(x, y)) +
11   geom_point() +
12   geom_line() +
13   geom_label(aes(x+.5,y+.5,label=coords))
```

R code Exa 16.3 Finding the probability of rolling a die and getting multiple faces

```
1 #Page Number: 292
2
3 trials <- 15
4 times <- 0:5
5 probab <- c(rep(1/6,6))
6 options(scipen = 999)
7 val <- round(dmultinom(times,size = trials,prob =
  probab),6)
8 cat(" Probability:" ,val)
```

R code Exa 16.4 Finding the probability of getting five 6s by rolling a die 15 times

```
1 #Page Number: 292
2
3 trials <- 15
4 times <- c(5,10)
5 probab <- c(rep(1/6,2))
6 res <- sum(round(dbinom(times,size = trials,prob =
  probab),4))
7 cat(" Probability:" ,res)
```

R code Exa 16.5 Testing the hypothesis of having 60 percent red marbles from a box having unknown composition of red and blue marbles

```
1 #Page Number: 293
2
3 n <- 15
4 X <- 11
5 prob <- 0.60
```

```

6 pcap <- round(X/n,4)
7 tstat <- round(sqrt(n)*(pcap-prob) / sqrt(prob*(1-
  prob)),4)
8 cat("Test Statistic:",tstat)
9
10 alpha <- 0.05
11 zt <- round(qnorm(alpha/2,lower.tail = F),2)
12 cat(tstat,"<",zt)

```

R code Exa 16.6 Determining the uniformity in composition of types of marble in a box

```

1 #Page Number: 294
2
3 n <- 60
4 x1 <- 22
5 x2 <- 17
6 x3 <- 21
7 k <- 3
8 p <- 1/3
9 np10 = np20 = np30 <- n*p
10
11 tstat <- round(((x1 - np10)^2 / np10) +((x2 - np20)
  ^2 / np20)+((x3 - np30)^2 / np30),3)
12 cat("Test Statistic:",tstat)
13
14 alpha <- 0.05
15 chi <- round(qchisq(alpha,df = k-1,lower.tail = F )
  ,2)
16 cat(tstat,"<",chi)

```

R code Exa 16.7 Testing the hypothesis of ratio of the offsprings in a biology experiment

```

1 #Page Number: 296
2
3 n <- 432
4 k <- 4
5 O1 <- 253
6 O2 <- 69
7 O3 <- 87
8 O4 <- 23
9
10 alpha <- 0.05
11
12 p1 <- 9/16
13 p2=p3 <- 3/16
14 p4 <- 1/16
15
16 E1 <- n*p1
17 E2 <- n*p2
18 E3 <- n*p3
19 E4 <- n*p4
20
21 tstat <- round(((O1-E1)^2 / E1)+((O2-E2)^2 / E2)+((
      O3-E3)^2 / E3)+((O4-E4)^2 / E4),4)
22 cat("Test Statistic:",tstat)
23
24 chi <- round(qchisq(alpha,k-1,lower.tail = F),2)
25 cat(tstat,"<",chi)

```

R code Exa 16.8 Goodness of fit test of survey data on a city daytime traffic

```

1 #Page Number: 297
2
3 data <- data.frame("Traffic Catagory" = c("PPV","LCV",
      "HCV"), "Sur 1" = c(436,391,297),"Sur 2" = c
      (520,401,319),"Sur 3" = c(376,281,191))

```

```

4 alpha <- 0.05
5 k <- 3
6
7 n1 <- sum(data$Sur.1)
8 n2 <- sum(data$Sur.2)
9 n3 <- sum(data$Sur.3)
10
11 p1 <- 0.40
12 p2 <- 0.35
13 p3 <- 0.25
14
15 O1 <- data$Sur.1
16 O2 <- data$Sur.2
17 O3 <- data$Sur.3
18
19
20 E1 <- n1*c(p1,p2,p3)
21 E2 <- n2*c(p1,p2,p3)
22 E3 <- n3*c(p1,p2,p3)
23
24 tstat1 <- sum(((O1 - E1)^2) /E1)
25 tstat2 <- sum(((O2 - E2)^2) /E2)
26 tstat3 <- sum(((O3 - E3)^2) /E3)
27 tot.tstat <- round(tstat1+tstat2+tstat3,4)
28 cat("Test Statistic:",tot.tstat)
29
30 chi <- round(qchisq(alpha,3*(k-1),lower.tail = F),3)
31 cat(tot.tstat,"<",chi)

```

Chapter 17

Hypothesis Testing to Compare Multiple Populations

R code Exa 17.1 Computing the population mean variance and applying 1 way anova test on experimental data of number of seconds taken to help in presence of other persons

```
1 #Page Number: 308
2
3 ze <- c(25,30,20,32,NA)
4 ze1 <- na.exclude(ze)
5 tw <- c(30,33,29,40,36)
6 fo <- c(32,39,35,41,44)
7 k <- 3
8 n1 <- length(ze1)
9 n2 <- length(tw)
10 n3 <- length(fo)
11
12 nT <- n1+n2+n3
13 s1 <- round(sd(ze, na.rm = T), 3)
14 s2 <- round(sd(tw), 3)
15 s3 <- round(sd(fo), 3)
16
17 sp <- round(sqrt((((n1-1)*(s1)^2)+((n2-1)*(s2)^2)+((
```

```

      n3-1)*(s3)^2)) / (nT-k)),3)
18 cat(" Pooled Sample SD:" ,sp)
19
20 c <- round(1+(1/(3*(k-1))) * ((1/(n1-1)+1/(n2-1)+1/(
      n3-1)) - (1/(nT-k))),4)
21 cat(" Constant:" ,c)
22
23 tstat <- round((2/c) *(((nT-k)*log(sp))-(((n1-1)*log
      (s1)) +((n2-1)*log(s2))+((n3-1)*log(s3))))),4)
24 cat(" Test Statistics:" ,tstat)
25
26 alpha <- 0.05
27 chi <- round(qchisq(alpha,(k),lower.tail = F),2)
28 cat(tstat,"<" ,chi)
29
30
31 xbar1 <- mean(ze1)
32 xbar2 <- mean(tw)
33 xbar3 <- mean(fo)
34 xbar <- round((n1*xbar1+n2*xbar2+n3*xbar3)/nT,4)
35 SSP <- round(n1*(xbar1-xbar)^2+n2*(xbar2-xbar)^2+n3*
      (xbar3-xbar)^2,2)
36 SSE <- round((nT - k)*(sp)^2,2)
37 TSS <- SSP + SSE
38 MSP <- SSP / (k-1)
39 MSE <- round(SSE / (nT-k),2)
40 Anova1 <- round(MSP / MSE,2)
41
42 tab <- data.frame("TSS"=TSS,"SSP"=SSP,"SSE"=SSE,"MSP
      "=MSP,"MSE"=MSE,"TStat"=Anova1)
43 print(tab)
44
45 ft <- qf(alpha,k-1,nT-k,lower.tail = F)
46 cat(Anova1,">" ,ft)

```

R code Exa 17.2 Classification of two way ANOVA and Barlett's test on data studying weight gains by rats influenced by the source of protein

```

1 #Page Number: 310
2
3 data <- data.frame("Diet-1" = c
  (90,94,86,63,81,89,63,82,63,81), "Diet-2" = c
  (87,70,95,71,75,84,62,72,81,93), "Diet-3" = c
  (83,82,85,63,72,85,64,80,82,83), "Diet-4" = c
  (107,102,102,93,111,128,56,97,80,103), "Diet-5" = c
  (96,72,76,70,79,89,70,91,63,102), "Diet-6" = c
  (111,100,102,93,101,104,72,92,87,112))
4
5 concat = c(t(as.matrix(data)))
6 f = c("Diet-1", "Diet-2", "Diet-3", "Diet-4", "Diet-5", "
  Diet-6")
7 k <- 6
8 n <- 10
9 data
10
11 pop <- gl(k, 1, n*k, factor(f))
12 block = gl(n, k, k*n)
13 anov2 = aov(concat ~ pop + block)
14 summary(anov2)
15
16
17 nT <- 60
18 s1 <- round(sd(data$Diet.1), 4)
19 s2 <- round(sd(data$Diet.2), 4)
20 s3 <- round(sd(data$Diet.3), 4)
21 s4 <- round(sd(data$Diet.4), 4)
22 s5 <- round(sd(data$Diet.5), 4)
23 s6 <- round(sd(data$Diet.6), 4)
24
25 sp <- round(sqrt((((n-1)*(s1)^2)+((n-1)*(s2)^2)+((n
  -1)*(s3)^2)+((n-1)*(s4)^2)+((n-1)*(s5)^2)+((n-1)*
  (s6)^2)) / (nT-k)), 4)
26 cat("Pooled Sample SD:", sp)

```

```

27
28 c <- round(1 + (1/(3*(k-1))) * (((1/(n-1))*6) - (1/(
      nT-k))),4)
29 cat(" Constant:" ,c)
30
31 tstat <- round((2/c) *(((nT-k)*log(sp)) - (((n-1)*
      log(s1)+(n-1)*log(s2)+(n-1)*log(s3)+(n-1)*log(s4)
      +(n-1)*log(s5)+(n-1)*log(s6))))),4)
32 cat(" Test Statistics:" ,tstat)
33
34 alpha <- 0.05
35 chi <- qchisq(alpha,(k),lower.tail = F)
36 cat(tstat,"<",round(chi,2))
37
38 #The answer may vary due to difference in
      representation.

```

Chapter 18

Quality Management Using Statistics

R code Exa 18.1 To test whether the manufacturing process is in control by construction of x chart and s chart from the dataset of JetMate

```
1 #Page Number: 336
2
3 library(qcc)
4
5 data <- read.csv("18_1.csv")
6 chart <- qcc.groups(data = data$JET, sample = data$
   GROUP)
7
8
9 plot.chart.R <- qcc(chart, type = "xbar", std.dev = "
   UWAVE-R")
10 summary(plot.chart.R)
11 plot.chart.SD <- qcc(chart, type = "xbar", std.dev = "
   UWAVE-SD")
12 summary(plot.chart.SD)
13
14
15 plot.chart_R <- qcc(chart, type = "R")
```

```

16 summary(plot.chart_R)
17
18 plot.chart_S <- qcc(chart,type = "S")
19 summary(plot.chart_S)
20 # The answer may slightly vary due to rounding off
    values.
21 # The answer may vary due to difference in
    representation.

```

R code Exa 18.2 To test whether the bottling process is in control by construction of x chart and r chart and s chart from the dataset of BottlePack company

```

1 #Page Number: 338
2
3 library(qcc)
4
5 data <- data.frame("Sam"=c(rep(1,5),rep(2,5),rep
    (3,5),rep(4,5),rep(5,5),rep(6,5),rep(7,5)), "Obs"=
    c
    (1.014,1.013,1.001,1.017,1.019,1.007,0.992,0.986,1.004,0.985,1.00
    )
6 data
7 chart <- qcc.groups(data = data$Obs, sample = data$
    Sam)
8
9
10 plot.chart_R <- qcc(chart,type = "R")
11 summary(plot.chart_R)
12
13 plot.chart.SD <- qcc(chart,type = "xbar",std.dev = "
    UWAVE-SD")
14 summary(plot.chart.SD)
15
16

```

```

17 plot.chart_S <- qcc(chart, type = "S")
18 summary(plot.chart_S)
19
20 #The answer may vary due to difference in
    representation.

```

R code Exa 18.3 To test whether the manufacturing process is in control by construction of s chart from the dataset of Blackstone company

```

1 #Page Number: 342
2
3 library(qcc)
4
5 data <- data.frame("Sam" = c(rep(1,3), rep(2,3), rep
    (3,3), rep(4,3), rep(5,3), rep(6,3), rep(7,3), rep
    (8,3), rep(9,3), rep(10,3)), "Obs" = c
    (40,33,30,34,21,28,27,32,36,18,28,24,40,40,38,30,20,18,30,28,31,30,
    )
6 chart <- qcc.groups(data = data$Obs, sample = data$
    Sam)
7
8 plot.chart_S <- qcc(chart, type = "S")
9 summary(plot.chart_S)

```

R code Exa 18.4 To test whether the manufacturing process is in control by construction of p chart from the dataset of ChipItGood company

```

1 #Page Number: 344
2
3 library(qcc)
4
5 data <- c
    (6,10,8,5,7,4,12,15,8,15,20,4,10,14,7,5,8,6)

```

```
6 n <- 100
7 plot.p <- qcc(data, type = "p", sizes = n)
8 summary(plot.p)
9
10 #The answer may vary due to difference in
    representation.
```
